

# *Glyceria notata*

## Habitus and growth type

Plant height [m]: **0.65**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **36.78**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.2**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **5.1**

Moisture indicator value: **9.7**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.13**

Disturbance frequency (herb layer): **1.63**

Disturbance severity: **0.39**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.21**

Grazing pressure: **0.16**

Soil disturbance: **0.12**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) Phragmito-Magnocaricetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q52 Small-helophyte bed](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Forest**

## **Distribution**

Continentality: **6**

Continentality amplitude: **5**