Gypsophila struthium

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9.3**
Moisture indicator value: **2.4**
Reaction indicator value: **7.9**
Nutrient indicator value: **3.5**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **1.91**
Disturbance frequency (herb layer): **1.86**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.11**
Grazing pressure: **0.3**
Soil disturbance: **0.21**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S65 Mediterranean gypsum scrub**
Constant species of EUNIS habitats: **S65 Mediterranean gypsum scrub**
Dominant species of EUNIS habitats: **S65 Mediterranean gypsum scrub**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz