

# *Herniaria hirsuta*

## Habitus and growth type

Plant height [m]: **0.07**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **21.23**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.51**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Moisture indicator value: **3.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **2.12**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.68**

Mowing frequency: **0.4**

Grazing pressure: **0.29**

Soil disturbance: **0.44**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S66 Mediterranean halo-nitrophilous scrub](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree),**

**Synanthropic**

## **Distribution**

Continentality: **6**

Continentality amplitude: **6**