

# *Hieracium virosum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.74**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **3.5**

Moisture indicator value: **5.3**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.65**

Grazing pressure: **0.38**

Soil disturbance: **0.17**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**