

Hierochloa odorata

Habitus and growth type

Plant height [m]: **0.46**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm²/mg]: **41.51**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.9**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.2**

Temperature indicator value: **4.8**

Moisture indicator value: **7.7**

Reaction indicator value: **4.8**

Nutrient indicator value: **2.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.79**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.44**

Grazing pressure: **0.19**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Grassland (non-alpine, non-saline)**

Distribution

Continentality: 7

Continentality amplitude: 6