Hippocrepis unisiliquosa

Habitus and growth type

Plant height [m]: **0.19**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **17.73**

Fruit, seed and dispersal

Seed mass [mg]: **5.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **11**
Moisture indicator value: **2.5**
Reaction indicator value: **8**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.62**
Disturbance frequency (herb layer): **1.9**
Disturbance severity: **0.79**
Disturbance severity (herb layer): **0.6**
Mowing frequency: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.25**
Soil disturbance: **0.37**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz