

# *Chamaemelum nobile*

## Habitus and growth type

Plant height [m]: **0.24**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.19**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Reaction indicator value: **4**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.97**

Disturbance frequency (herb layer): **2.29**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.66**

Grazing pressure: **0.59**

Soil disturbance: **0.33**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#)

Constant species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**