

Chenopodium ficifolium

Habitus and growth type

Plant height [m]: **0.65**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **18.92**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.37**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.1**

Moisture indicator value: **6**

Reaction indicator value: **6**

Nutrient indicator value: **7.3**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.84**

Disturbance frequency (herb layer): **1.93**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.81**

Mowing frequency: **0.53**

Grazing pressure: **0.17**

Soil disturbance: **0.66**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QI \(BID\) Bidentetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA221 Atlantic saltmarsh driftline](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Synanthropic**

Distribution

Continentality: 7

Continentality amplitude: 7