# *Cherleria baldaccii*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Disturbance indicator values

Disturbance frequency: **0.22**
Disturbance frequency (herb layer): **1.11**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.2**
Mowing frequency: **0.04**
Grazing pressure: **0.21**
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CU (SES) *Elyno-Seslerietea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **T39 Mediterranean and Balkan subalpine *Pinus heldreichii-Pinus peuce* forest**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*