

Laserpitium latifolium

Habitus and growth type

Plant height [m]: **1.15**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **14.22**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **8.63**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **4.8**

Moisture indicator value: **4.5**

Reaction indicator value: **7.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.33**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.17**

Grazing pressure: **0.22**

Soil disturbance: **0.11**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [R51 Thermophilous forest fringe of base-rich soils](#), [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#), [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **3**