Lysimachia vulgaris

Habitus and growth type

Plant height [m]: **0.84**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **26.37**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.28**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**
Temperature indicator value: **5.1**
Moisture indicator value: **8.6**
Reaction indicator value: **6.3**
Nutrient indicator value: **5.3**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.64**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.33**
Grazing pressure: **0.17**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea*, OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q53 Tall-sedge bed, R35 Moist or wet mesotrophic to eutrophic hay meadow, S92 Salix fen scrub, T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils, T15 Broadleaved swamp forest on non-acid peat**

Constant species of EUNIS habitats: **Q22 Poor fen, Q24 Intermediate fen and soft-water spring mire, Q42 Extremely rich moss-sedge fen, Q51 Tall-helophyte bed, Q53 Tall-sedge bed, R35 Moist or wet mesotrophic to eutrophic hay meadow, R37 Temperate and boreal moist or wet oligotrophic grassland, R55 Lowland moist or wet tall-herb and fern fringe, S92 Salix fen scrub, T11 Temperate Salix and Populus riparian forest, T12 Alnus glutinosa-Alnus incana forest on riparian and mineral soils, T13 Temperate hardwood riparian forest, T15 Broadleaved swamp forest on non-acid peat, T16 Broadleaved mire forest on acid peat**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz