Meniocus linifolius

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.73**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.8**
Moisture indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **2.11**
Disturbance frequency (herb layer): **2.07**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.28**
Grazing pressure: **0.41**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R15 Continental dry rocky steppic grassland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

and dwarf scrub on chalk outcrops, **R1C Desert steppe**, **S67 Aralo-Caspian semi-desert**

Constant species of EUNIS habitats: **R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops**, **R1C Desert steppe**, **S67 Aralo-Caspian semi-desert**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz