

Minuartia viscosa

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **6.7**

Moisture indicator value: **2.9**

Reaction indicator value: **4**

Nutrient indicator value: **1.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.08**

Disturbance frequency (herb layer): **2.08**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.46**

Grazing pressure: **0.37**

Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) *Koelerio-Coryneporetea canescentis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R11 Pannonian and Pontic sandy steppe](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline)**

Distribution

Continentality: 7

Continentality amplitude: 4