

Muscari botryoides

Habitus and growth type

Plant height [m]: **0.17**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **28.94**

Flower

Flowering period: **January-April**

Fruit, seed and dispersal

Seed mass [mg]: **2.65**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**

Temperature indicator value: **5.9**

Moisture indicator value: **4.9**

Reaction indicator value: **7.3**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.38**

Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.57**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.21**
Grazing pressure: **0.28**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **3**