Nananthea perpusilla

Habitus and growth type

Plant height [m]: **0.02**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **January-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **10**
Moisture indicator value: **5**
Reaction indicator value: **6**
Nutrient indicator value: **1**
Salinity indicator value: **3**

Disturbance indicator values

Disturbance frequency: **2.06**
Disturbance frequency (herb layer): **2.06**
Disturbance severity: **0.25**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.04**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.05**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz