

# *Nonea pulla*

## Habitus and growth type

Plant height [m]: **0.35**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **22.97**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **7.28**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.6**

Moisture indicator value: **3.6**

Reaction indicator value: **7.8**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.76**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.89**

Grazing pressure: **0.32**

Soil disturbance: **0.22**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QB \(SIS\) Sisymbrietea](#), [QC \(CHE\) Chenopodietea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [R1B Continental dry grassland \(true steppe\)](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

## **Distribution**

Continentality: 7

Continentality amplitude: 5