

Odontarrhena borzaeana

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **3.8**

Reaction indicator value: **8**

Nutrient indicator value: **6**

Salinity indicator value: **2**

Disturbance indicator values

Disturbance frequency: **1.6**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.05**

Grazing pressure: **0.21**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **[CK \(SED\) Sedo-Scleranthetea](#)**

EUNIS habitat

Diagnostic species of EUNIS habitats: [N17 Black Sea coastal dune grassland \(grey dune\)](#)

Constant species of EUNIS habitats: [N17 Black Sea coastal dune grassland \(grey dune\)](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline)**