Ochlopoa infirma

Habitus and growth type

Plant height [m]: **0.05**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-September**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**
Temperature indicator value: **7.8**
Reaction indicator value: **5.8**
Nutrient indicator value: **5.2**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.35**
Disturbance frequency (herb layer): **1.84**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.67**
Mowing frequency: **0.41**
Grazing pressure: **0.32**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.38**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QE (POL) *Polygono-Poetea annuae***

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz