Orobanche artemisiae-campestris

Habitus and growth type

Plant height [m]: **0.33**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**
Temperature indicator value: **7.2**
Moisture indicator value: **3.9**
Reaction indicator value: **7.4**
Nutrient indicator value: **4.1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.09**
Disturbance frequency (herb layer): **1.63**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.48**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.38**
Grazing pressure: **0.2**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **FC (LAV) *Cisto-Lavanduletea stoechadis***, **QF (ART) *Artemisietea vulgaris***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz