

# *Paederota lutea*

## Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Temperature indicator value: **3.3**

Moisture indicator value: **7**

Reaction indicator value: **8**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.21**

Disturbance frequency (herb layer): **0.8**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.16**

Mowing frequency: **0**

Grazing pressure: **0.19**

Soil disturbance: **0.08**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) \*Asplenietea trichomanis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [U36 Temperate high-mountain base-rich inland cliff](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**