

# *Parentucellia viscosa*

## Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.21**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.02**

Dispersal mode: **Anemochory**

## Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Reaction indicator value: **6**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **1.18**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.64**

Mowing frequency: **0.44**

Grazing pressure: **0.26**

Soil disturbance: **0.34**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**