

# *Paronychia macrosepala*

## Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7.9**

Moisture indicator value: **3.7**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.26**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.11**

Grazing pressure: **0.29**

Soil disturbance: **0.28**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) Ononido-Rosmarinetea**, **FJ (TRA) Stipo-Trachynietea distachyae**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**