Paronychia suffruticosa

Habitus and growth type

Plant height [m]: **0.25**
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **7.08**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **8.6**
Moisture indicator value: **2.9**
Reaction indicator value: **6.6**
Nutrient indicator value: **3.4**
Salinity indicator value: **0.9**

Disturbance indicator values

Disturbance frequency: **0.32**
Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.21**
Mowing frequency: **0.06**
Grazing pressure: **0.42**
Soil disturbance: **0.15**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz