

# *Paspalum distichum*

## Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **1.45**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **8.5**

Moisture indicator value: **8.7**

Reaction indicator value: **6.7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.35**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.55**

Mowing frequency: **0.28**

Grazing pressure: **0.2**

Soil disturbance: **0.32**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [V14 Inundated or inundatable cropland, including rice fields](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Synanthropic**