

Poa palustris

Habitus and growth type

Plant height [m]: **0.74**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **28**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**

Temperature indicator value: **5.1**

Moisture indicator value: **8.4**

Reaction indicator value: **7**

Nutrient indicator value: **6.7**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.94**

Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.48**
Grazing pressure: **0.19**
Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[OD \(PHR\) *Phragmito-Magnocaricetea*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: 7
Continentality amplitude: 7