

Poa timoleontis

Habitus and growth type

Plant height [m]: **0.16**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **8.1**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.7**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.14**

Grazing pressure: **0.35**

Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FR \(DAP\) Daphno-Festucetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**