

Reynoutria japonica

Habitus and growth type

Plant height [m]: **1.59**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **18.92**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **2.39**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **6.2**

Moisture indicator value: **7.3**

Reaction indicator value: **5.3**

Nutrient indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.68**

Disturbance frequency (herb layer): **1.48**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.41**

Grazing pressure: **0.23**

Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QG \(EPI\) Epilobietea angustifolii](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V39 Mesic perennial anthropogenic herbaceous vegetation](#)

Dominant species of EUNIS habitats: [V39 Mesic perennial anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Synanthropic**

Distribution

Continentality: **4**

Continentality amplitude: **4**