

Rhaponticum coniferum

Habitus and growth type

Plant height [m]: **0.11**
Life span: **Perennial**
Life form: **Hemicryptophyte**
Spinescence: **spinescent**

Leaf

Specific leaf area [mm²/mg]: **10.13**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**
Ellenberg-type indicator values
Light indicator value: **9**
Moisture indicator value: **3.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**
Disturbance indicator values
Disturbance frequency: **0.37**
Disturbance frequency (herb layer): **1.55**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.14**

Grazing pressure: **0.35**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#), [FF \(LYG\) Lygeo sparti-Stipetea tenacissimae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [S61 Western basiphilous garrigue](#)

Constant species of EUNIS habitats: [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#), [S61 Western basiphilous garrigue](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**