Romulea revelieri

Habitus and growth type

Plant height [m]: **0.11**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9**
Moisture indicator value: **4.5**
Reaction indicator value: **3.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.06**
Disturbance frequency (herb layer): **2.06**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.55**
Mowing frequency: **0.02**
Grazing pressure: **0.35**
Soil disturbance: **0.3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz