

Rumex hydrolapathum

Habitus and growth type

Plant height [m]: **1.54**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **18.79**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **6.3**

Moisture indicator value: **9.7**

Reaction indicator value: **7**

Nutrient indicator value: **6.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.28**

Disturbance frequency (herb layer): **1.69**
Disturbance severity: **0.3**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.41**
Grazing pressure: **0.12**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q51 Tall-helophyte bed](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline)**

Distribution

Continentality amplitude: **5**