

Sagina maritima

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **14.3**

Flower

Flowering period: **January-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.14**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **6.3**

Reaction indicator value: **7.5**

Salinity indicator value: **3.8**

Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.11**

Grazing pressure: **0.36**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JA \(SAG\) *Saginetea maritimae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#), [MA251 Mediterranean upper saltmarsh](#), [N31 Atlantic and Baltic rocky sea cliff and shore](#)

Constant species of EUNIS habitats: [MA222 Atlantic upper saltmarsh](#), [MA251 Mediterranean upper saltmarsh](#), [N31 Atlantic and Baltic rocky sea cliff and shore](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal cliff, Saline vegetation, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**