

Sanguisorba officinalis

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.03**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **3.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **4.7**

Moisture indicator value: **6.4**

Reaction indicator value: **6.3**

Nutrient indicator value: **4**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.55**

Disturbance frequency (herb layer): **1.91**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **1.15**
Grazing pressure: **0.18**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#), [CM \(MOL\) *Molinio-Arrhenatheretea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [T3H Larix light taiga](#)

Constant species of EUNIS habitats: [R35 Moist or wet mesotrophic to eutrophic hay meadow](#), [R37 Temperate and boreal moist or wet oligotrophic grassland](#), [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Grassland (non-alpine, non-saline), Forest**

Distribution

Continentality: **6**
Continentality amplitude: **6**