

Saxifraga bryoides

Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Chamaephyte**

Leaf

Specific leaf area [mm²/mg]: **18.2**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **1.5**

Moisture indicator value: **5.4**

Reaction indicator value: **3**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.51**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0**
Grazing pressure: **0.19**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U22 Temperate high-mountain siliceous scree](#), [U32 Temperate high-mountain siliceous inland cliff](#)

Constant species of EUNIS habitats: [U22 Temperate high-mountain siliceous scree](#), [U32 Temperate high-mountain siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **3**