Saxifraga rotundifolia

Habitus and growth type

Plant height [m]: **0.35**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **3.6**
Moisture indicator value: **6**
Reaction indicator value: **7.6**
Nutrient indicator value: **5.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.11**
Disturbance frequency (herb layer): **1.11**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.01**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.18**
Soil disturbance: **0.09**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R56 Montane to subalpine moist or wet tall-herb and fern fringe**, **S25 Subalpine and subarctic deciduous scrub**
Constant species of EUNIS habitats: **R56 Montane to subalpine moist or wet tall-herb and fern fringe**, **S25 Subalpine and subarctic deciduous scrub**

Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz