# *Saxifraga squarrosa*

## Habitus and growth type

Plant height [m]: **0.04**
Life span: **Perennial**
Life form: **Chamaephyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Seed mass [mg]: **0.05**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **3**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **1**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.73**
Disturbance frequency (herb layer): **0.82**
Disturbance severity: **0.22**
Disturbance severity (herb layer): **0.19**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0**
Grazing pressure: **0.14**
Soil disturbance: **0.05**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **KC (ASP) *Asplenietea trichomanis***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **U36 Temperate high-mountain base-rich inland cliff**

Constant species of EUNIS habitats: **U36 Temperate high-mountain base-rich inland cliff**

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*