

# *Scorzoneroides helvetica*

## Habitus and growth type

Plant height [m]: **0.13**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.85**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.5**

Moisture indicator value: **4.8**

Reaction indicator value: **3**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1**

Disturbance frequency (herb layer): **1.45**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.04**

Grazing pressure: **0.25**

Soil disturbance: **0.09**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CT \(TRI\) \*Juncetea trifidi\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R43 Temperate acidophilous alpine grassland](#),  
[S22 Alpine and subalpine ericoid heath](#)

Constant species of EUNIS habitats: [R43 Temperate acidophilous alpine grassland](#),  
[S22 Alpine and subalpine ericoid heath](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Heathland, Scrub**

## Distribution

Continentality: **4**

Continentality amplitude: **2**