

# *Seseli libanotis*

## Habitus and growth type

Plant height [m]: **0.64**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.59**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **1.53**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **5.3**

Moisture indicator value: **3.2**

Reaction indicator value: **7.6**

Nutrient indicator value: **2.1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.9**

Disturbance frequency (herb layer): **1.63**  
Disturbance severity: **0.49**  
Disturbance severity (herb layer): **0.42**  
Mowing frequency: **0.44**  
Grazing pressure: **0.26**  
Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) \*Brachypodio pinnati-Betuletea pendulae\*](#), [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#), [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: 7