

Silene conica

Habitus and growth type

Plant height [m]: **0.17**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **15.89**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.33**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **7.6**

Moisture indicator value: **1.8**

Nutrient indicator value: **2.5**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.29**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.31**

Grazing pressure: **0.29**

Soil disturbance: **0.4**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) *Sedo-Scleranthetea*](#), [EJ \(TRA\) *Stipo-Trachynietea distachyae*](#), [KH \(DRY\) *Drypidetea spinosae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R11 Pannonian and Pontic sandy steppe](#), [S34 Balkan-Anatolian submontane genistoid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **6**