Spiraea crenata

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **5.1**
Moisture indicator value: **4.6**

Disturbance indicator values

Disturbance frequency: **1.38**
Disturbance frequency (herb layer): **1.56**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.53**
Grazing pressure: **0.37**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **CD (RHA) *Crataego-Prunetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S36 Low steppic scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz