

# *Stachys cretica*

## Habitus and growth type

Plant height [m]: **0.51**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **3.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **3.6**

Reaction indicator value: **7.7**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.67**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.36**

Grazing pressure: **0.3**

Soil disturbance: **0.22**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [T38 Mediterranean montane Cedrus forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**