

Stachys scardica

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **6.6**

Temperature indicator value: **6.4**

Moisture indicator value: **4.9**

Reaction indicator value: **5.9**

Nutrient indicator value: **3.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.24**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.14**

Grazing pressure: **0.24**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#), [CO \(ERI\) Erico-Pinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**