Stuckenia filiformis

Habitus and growth type

Plant height [m]: **0.35**
Life span: **Perennial**
Life form: **Hydrophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.11**
Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**
Temperature indicator value: **4.6**
Moisture indicator value: **11.8**
Reaction indicator value: **5.2**
Nutrient indicator value: **3.4**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.61**
Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.25**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.01**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.02**
Soil disturbance: **0.02**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **NB (POT) *Potamogetonetea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

Distribution

Continentality: **7**
Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz