

Suaeda acuminata

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.43**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **6.5**

Moisture indicator value: **6.3**

Disturbance indicator values

Disturbance frequency: **2.1**

Disturbance frequency (herb layer): **2.12**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.08**

Grazing pressure: **0.45**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DB \(FEP\) Festuco-Puccinellietea](#), [ME \(THE\) Thero-Salicornietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R64 Semi-desert salt pan](#), [S94 Semi-desert](#)

[riparian scrub](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Saline vegetation, Scrub**