# *Tanacetum balsamitoides*

## Habitus and growth type

Plant height [m]: **0.7**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6**
Moisture indicator value: **5**
Reaction indicator value: **6**
Nutrient indicator value: **7**
Salinity indicator value: **0**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***

### Broad habitat

Occurrence in broad habitats: **Synanthropic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*