

# *Taraxacum stevenii*

## Habitus and growth type

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **28.8**

## Fruit, seed and dispersal

Seed mass [mg]: **1**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Disturbance indicator values

Disturbance frequency: **1.71**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.25**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.01**

Grazing pressure: **0.18**

Soil disturbance: **0.1**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KF \(LAM\) \*Lamio tomentosii-Chaerophylletea humilis\*](#), [LA \(HER\) \*Salicetea herbaceae\*](#)

Broad habitat

Occurrence in broad habitats: **Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**