

# *Tolpis barbata*

## Habitus and growth type

Plant height [m]: **0.5**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **32.44**

## Fruit, seed and dispersal

Seed mass [mg]: **0.16**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **2**

Reaction indicator value: **4.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.16**

Disturbance frequency (herb layer): **2**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.53**

Grazing pressure: **0.32**

Soil disturbance: **0.23**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

Constant species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**