

# *Trifolium striatum*

## Habitus and growth type

Plant height [m]: **0.19**

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **20.94**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Seed mass [mg]: **1.64**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.3**

Moisture indicator value: **2.5**

Reaction indicator value: **3.7**

Nutrient indicator value: **2.2**

Salinity indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **1.81**

Disturbance frequency (herb layer): **2.17**  
Disturbance severity: **0.54**  
Disturbance severity (herb layer): **0.54**  
Mowing frequency: **0.67**  
Grazing pressure: **0.35**  
Soil disturbance: **0.33**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Coryneporetea canescentis\*](#), [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#), [R34 Submediterranean moist meadow](#)

Constant species of EUNIS habitats: [R24 Iberian summer pasture \(vallicar\)](#), [R34 Submediterranean moist meadow](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**  
Continentality amplitude: **4**