Trichophorum alpinum

Habitus and growth type

Plant height [m]: **0.17**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **9.43**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.32**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **4**
Moisture indicator value: **9.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.14**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.18**
Mowing frequency: **0.39**
Grazing pressure: **0.1**
Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **PA (SCH) *Scheuchzerio palustris-Caricetea fuscae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q42 Extremely rich moss-sedge fen**, **Q44 Calcareous quaking mire**, **Q45 Arctic-alpine rich fen**
Constant species of EUNIS habitats: **Q42 Extremely rich moss-sedge fen**, **Q44 Calcareous quaking mire**

Broad habitat

Occurrence in broad habitats: **Spring, Mire**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz