

# *Trisetaria ovata*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **1**

Reaction indicator value: **3**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.83**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.59**

Grazing pressure: **0.3**

Soil disturbance: **0.25**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**