Vaccinium macrocarpon

Habitus and growth type

Plant height [m]: **0.6**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm^2/mg]: **12.59**

Fruit, seed and dispersal

Seed mass [mg]: **0.79**
Dispersal mode: **Endozoochory, Anthropochory**
Dispersal distance class: **6, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Disturbance indicator values

Disturbance frequency: **1.14**
Disturbance frequency (herb layer): **1.25**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.01**
Grazing pressure: **0.35**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **PB (OXY) *Oxycocco-Sphagnetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1A Atlantic and Baltic coastal dune scrub**

Broad habitat

Occurrence in broad habitats: **Mire**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Distribution

Continentality: **5**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz